## For Research Use Only

## CD46 Polyclonal antibody

Catalog Number: 28936-1-AP



**Basic Information** 

Catalog Number:

Size:

GenBank Accession Number: BC030594

28936-1-AP

GeneID (NCBI):

150ul , Concentration: 340 ug/ml by

Nanodrop: **UNIPROT ID:** P15529

Rabbit Full Name:

Isotype: CD46 molecule, complement

IgG regulatory protein Immunogen Catalog Number: Calculated MW: AG30376 384 aa. 43 kDa

> Observed MW: 50-70 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Purification Method:** Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:50-1:500 IF/ICC 1:200-1:800

Positive Controls:

WB: HeLa cells, Jurkat cells, A549 cells

IHC: human colon cancer tissue,

IF/ICC: HeLa cells,

## **Background Information**

CD46, also known as Membrane cofactor protein (MCP) or MIC10, is a ubiquitously expressed type type-I transmembrane glycoprotein that acts as a cofactor for complement factor I to mediate inactivation of C3b and C4b deposited on host cells. CD46 is a receptor for a large number of pathogens, including measles virus and adenovirus (PMID: 20941397). CD46 plays a role in linking innate and adaptive immune responses. It may be involved in the fusion of the spermatozoa with the oocyte during fertilization. Mutations of the CD46 gene have been associated with some diseases including hemolytic uremic syndrome (PMID: 26054645). CD46 is expressed on most cells as four isoforms that are derived via alternative splicing (PMID: 15324743).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

E: proteintech@ptglab.com


W: ptglab.com

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




HeLa cells were subjected to SDS PAGE followed by western blot with 28936-1-AP (CD46 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 28936-1-AP (CD46 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 28936-1-AP (CD46 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using CD46 antibody (28936-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).