For Research Use Only

Phospho-p70(S6K) (Thr229) Polyclonal antibody

www.ptglab.com

Purification Method:

WB 1:1000-1:6000 IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Catalog Number: 28988-1-AP

Basic Information

Catalog Number:

28988-1-AP BC053365 GeneID (NCBI):

100ul, Concentration: 800 ug/ml by 6198

Nanodrop: **UNIPROT ID:** Source: P23443

Isotype: ribosomal protein S6 kinase, 70kDa,

polypeptide 1 IgG

Calculated MW: 59 kDa Observed MW: 65-85 kDa

Full Name:

GenBank Accession Number:

Applications

Tested Applications: WB, IF/ICC, ELISA

Species Specificity:

Rabbit

Positive Controls:

WB: λ phosphatase treated HeLa cells, IF/ICC: λ phosphatase treated HeLa cells,

Background Information

The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K), which is a serine/threonine kinase $regulated\ by\ phosphoinositide\ 3-kinase\ (PI3K)/mammalian\ target\ of\ rapamycin\ (mTOR)\ pathway.\ P70S6K\ plays\ a$ crucial role in controlling cell cycle, growth and survival. The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation and metabolism and is the central regulator of translation of some components of protein synthesis system. Due to alternative translation two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1). (PMID: 25100792, PMID: 24970012, PMID: 21602892)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HeLa, phosphatase inhibitor treated and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 28988-1-AP (Phospho-p70(S6f) (Thr229) antibody) at dilution of 1:3000 incubated at room temperature for 1 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Immunofluorescent analysis of (4% PFA) fixed λ phosphatase treated HeLa cells using Phosphop70(56K) (Thr229) antibody (28988-1-AP) at dilution of 1:200 and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).