For Research Use Only

Phospho-eEF2K (Ser366) Polyclonal antibody

Proteintech

Antibodies I ELISA kits I Proteins

www.ptglab.com

Catalog Number: 29032-1-AP

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

29032-1-AP

GeneID (NCBI):

iningerialilinity purification

100ul , Concentration: 300 ug/ml by

29904

BC032665

Recommended Dilutions: WB 1:1000-1:4000

Nanodrop; Source:

UNIPROT ID:

000418 Full Name:

Rabbit Fullsotype: eu

eukaryotic elongation factor-2 kinase

IgG Calculated MW:

725 aa, 100 kDa Observed MW:

100 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Positive Controls:

WB: Jurkat cells, λ phosphatase treated Jurkat cells

Background Information

Eukaryotic Elongation Factor-2 Kinase (eEF2K) acts as a negative regulator of protein synthesis, translation, and cell growth. As a structurally unique member of the alpha-kinase family, eEF2K is essential to cell survival under stressful conditions, as it contributes to both cell viability and proliferation. eEF2K is regulated by various mechanisms, including phosphorylation through residues and autophosphorylation. eEF2K is regulated by various mechanisms, including phosphorylation through residues and autophosphorylation. eEF2K is downregulated through the phosphorylation of multiple sites via mTOR signaling and upregulated via the AMPK pathway. In the S6K-mediated pathway, RPS6KB1 or p70S6K, can phosphorylate eEF2K on Ser-366, rendering it inactive(PMID: 11500364, PMID: 34532346). The calculated molecular weight of eEF2K is 82kDa, and probably due to phosphorylation modification, resulting in a larger molecular weight, around 100kDa.

Storage

Storage:

Store at -20°C.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated and λ phosphatase treated Jurkat cells were subjected to SDS PAGE followed by western blot with 29032-1-AP (Phospho-eEF2K (Ser366) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.