

For Research Use Only

Phospho-PRAS40 (Thr246) Polyclonal antibody



Catalog Number: 29072-1-AP

Basic Information

Catalog Number: 29072-1-AP	GenBank Accession Number: BC051844	Purification Method: Antigen affinity purification
Size: 100ul , Concentration: 240 µg/ml by Nanodrop;	GeneID (NCBI): 84335	Recommended Dilutions: WB 1:2000-1:10000
Source: Rabbit	Full Name: AKT1 substrate 1 (proline-rich)	
Isotype: IgG	Calculated MW: 27 kDa	
	Observed MW: 40 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : IGF-1 treated MCF-7 cells, Calyculin A treated PC-3 cells
Species Specificity: Human	

Background Information

PRAS40 is expressed in a variety of tissues in vivo and has multiple phosphorylation sites, which its activity is closely related to phosphorylation. Studies have shown that PRAS40 is involved in regulating cell growth, cell apoptosis, oxidative stress, autophagy and angiogenesis, as well as various of signalling pathways such as mammalian target of mammalian target rapamycin (mTOR), protein kinase B (PKB/Akt), nuclear factor kappa-B (NF-κB), proto-oncogene serine/threonine-protein kinase PIM-1 (PIM1) and pyruvate kinase M2 (PKM2). PGK1 phosphorylated PRAS40 at Threonine 246, which could be inhibited by blocking the interaction. PRAS40 can be phosphorylated by p-AKT at Thr246 site and promote the signal downstream to the mTOR signaling pathway. (PMID: 35058442, PMID: 32855714)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated MCF-7 cells and IGF-1 treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 29072-1-AP (Phospho-PRAS40 (Thr246) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Non-treated PC-3 cells and Calyculin A treated PC-3 cells were subjected to SDS PAGE followed by western blot with 29072-1-AP (Phospho-PRAS40 (Thr246) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.