

For Research Use Only

# SULT2B1 Polyclonal antibody

Catalog Number: 29185-1-AP



## Basic Information

<b>Catalog Number:</b> 29185-1-AP	<b>GenBank Accession Number:</b> BC034694	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 1200 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 6820	<b>Recommended Dilutions:</b> WB 1:1000-1:4000
<b>Source:</b> Rabbit	<b>Full Name:</b> sulfotransferase family, cytosolic, 2B, member 1	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 365 aa, 41 kDa	
<b>Immunogen Catalog Number:</b> AG30209	<b>Observed MW:</b> 41 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : A549 cells, mouse skin tissue, LNCaP cells, MCF-7 cells
<b>Species Specificity:</b> Human, Mouse	

## Background Information

SULT2B1 is a member of hydroxysteroid sulfotransferase (SULT) family that catalyzes sulfonation of hormones and other molecules. SULT2B1 encodes two isoforms, SULT2B1a and SULT2B1b, that differ only at their amino termini but have distinct biochemical properties as well as tissue distribution. The human fetal brain appears to only express the SULT2B1a isoform, while SULT2B1b is the predominant hydroxysteroid sulfotransferase expressed in human skin. Northern blot analysis revealed SULT2B1a has a limited tissue distribution in liver, adrenal and stomach, whereas SULT2B1b can be detected in a variety of hormone-responsive tissues including placenta, ovary, uterus, and prostate.

## Storage


**Storage:**  
Store at -20°C.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29185-1-AP (SULT2B1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.