

For Research Use Only

Phospho-p70(S6K) (Thr421/Ser424) Polyclonal antibody



Catalog Number: 29248-1-AP

Basic Information

Catalog Number: 29248-1-AP	GenBank Accession Number: BC053365	Purification Method: Antigen affinity purification
Size: 100ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 6198	Recommended Dilutions: WB 1:1000-1:4000 IF 1:50-1:500
Source: Rabbit	Full Name: ribosomal protein S6 kinase, 70kDa, polypeptide 1	
Isotype: IgG	Calculated MW: 59 kDa	
	Observed MW: 65-85 kDa	

Applications

Tested Applications: IF, WB, ELISA	Positive Controls: WB : λ phosphatase treated MCF-7 cells, IF : λ phosphatase treated HeLa cells,
Species Specificity: Human	

Background Information

The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K), which is a serine/threonine kinase regulated by phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin (mTOR) pathway. P70S6K plays a crucial role in controlling cell cycle, growth and survival. The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation and metabolism and is the central regulator of translation of some components of protein synthesis system. Due to alternative translation two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1). In the conventional model, Thr421/Ser424 sites are phosphorylated before Thr389 phosphorylation by mammalian target of rapamycin complex 1 (mTORC1) and subsequent Thr229 phosphorylation by PDK1. (PMID: 25100792, PMID: 24970012, PMID: 21602892)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated MCF-7 cells, phosphatase inhibitor treated and λ phosphatase MCF-7 cells were subjected to SDS PAGE followed by western blot with 29248-1-AP (Phospho-p70(S6K) (Thr421/Ser424) antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Immunofluorescent analysis of (4% PFA) fixed λ phosphatase treated HeLa cells using Phospho-p70(S6K) (Thr421/Ser424) antibody (29248-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).