For Research Use Only

STK11/LKB1 Polyclonal antibody Catalog Number:29323-1-AP Featured Product 1 Publication

Featured Product 1 Publications


oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 29323-1-AP	GenBank Accession Number: BC007981 GeneID (NCBI): 6794 UNIPROT ID: Q15831 Full Name: serine/threonine kinase 11 Calculated MW: 49 kDa Observed MW: 50-55 kDa		Purification Method: Antigen affinity purification		
	Size: 150ul, Concentration: 450 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG30903			Recommended Dilutions: WB 1:2000-1:10000 IHC 1:300-1:1200		
Applications	Tested Applications:		Positive Cor	sitive Controls:		
	WB, IHC, ELISA Cited Applications: WB	WB: A431 Cetts, MCF-7 Cetts, HER-2931 ted Applications: IHC : human lung cancer tissue, human tissue, mouse testis tissue, human lym				
	Species Specificity: Human, Mouse Cited Species:					
	mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
	STK11(serine/threonine-protein kinase 11) is also named as LKB1, PJS, and belongs to the protein kinase superfamily. It controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a rol in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response. The tumour suppressor protein LKB1 is a serine/threonine kinase that has been causally linked to Peutz-Jeghers syndrome (PJS) Defects in STK11 are a cause of Peutz-Jeghers syndrome (PJS) and defects in STK11 have been associated with testicular germ cell tumor (TGCT) and some sporadic cancers, especially lung cancers. STK11 has 2 isoforms with MW of 49 kDa and 45 kDa, and can be detected as 50-54 kDa after posttranslational modification.					
Background Information	superfamily. It controls the activity o in various processes such as cell met suppressor protein LKB1 is a serine/th Defects in STK11 are a cause of Peutz testicular germ cell tumor (TGCT) and	of AMP-activated prot abolism, cell polarit preonine kinase that -Jeghers syndrome (d some sporadic can	tein kinase (AMP y, apoptosis and has been causal PJS) and defects cers, especially l	K) family members, thereby playing a ro DNA damage response. The tumour ly linked to Peutz-Jeghers syndrome (PJS in STK11 have been associated with ung cancers. STK11 has 2 isoforms with		
Background Information	superfamily. It controls the activity of in various processes such as cell met suppressor protein LKB1 is a serine/th Defects in STK11 are a cause of Peutz testicular germ cell tumor (TGCT) and MW of 49 kDa and 45 kDa, and can be	of AMP-activated prof abolism, cell polarit preonine kinase that -Jeghers syndrome (d some sporadic canc e detected as 50-54 k	tein kinase (AMP y, apoptosis and has been causal PJS) and defects cers, especially l Da after posttrar	K) family members, thereby playing a rol DNA damage response. The tumour Ly linked to Peutz-Jeghers syndrome (PJS in STK11 have been associated with ung cancers. STK11 has 2 isoforms with slational modification.		
Background Information	superfamily. It controls the activity of in various processes such as cell met suppressor protein LKB1 is a serine/th Defects in STK11 are a cause of Peutz testicular germ cell tumor (TGCT) and MW of 49 kDa and 45 kDa, and can be Author Put	of AMP-activated prof abolism, cell polarit reconine kinase that -Jeghers syndrome (d some sporadic can e detected as 50-54 k	tein kinase (AMP y, apoptosis and has been causal PJS) and defects cers, especially l	K) family members, thereby playing a ro DNA damage response. The tumour ly linked to Peutz-Jeghers syndrome (PJS in STK11 have been associated with ung cancers. STK11 has 2 isoforms with		
	superfamily. It controls the activity of in various processes such as cell met suppressor protein LKB1 is a serine/th Defects in STK11 are a cause of Peutz testicular germ cell tumor (TGCT) and MW of 49 kDa and 45 kDa, and can be Author Put	of AMP-activated prof abolism, cell polarit preonine kinase that -Jeghers syndrome (d some sporadic can e detected as 50-54 k comed ID Jou 731709 Free ter shipment.	tein kinase (AMP y, apoptosis and has been causal PJS) and defects cers, especially l Da after posttran	K) family members, thereby playing a ro DNA damage response. The tumour Ly linked to Peutz-Jeghers syndrome (PJS in STK11 have been associated with ung cancers. STK11 has 2 isoforms with slational modification. Application		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 29323-1-AP (STK11 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.





WB result of STK11/LKB1 antibody (29323-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STK11/LKB1 transfected MCF-7 cells. WB result of STK11/LKB1 antibody (29323-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STK11/LKB1 transfected A431 cells.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 29323-1-AP (STK11/LKB1 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).