## For Research Use Only

## Annexin A4 Polyclonal antibody

Catalog Number: 29508-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

29508-1-AP BC000182 GeneID (NCBI):

150ul, Concentration: 650 ug/ml by

Nanodrop: **UNIPROT ID:** P09525 Rabbit Full Name: Isotype: annexin A4 IgG Calculated MW: Immunogen Catalog Number: 36 kDa

AG30827 Observed MW:

36 kDa

**Applications Tested Applications:** 

WB, ELISA

Species Specificity: Human, rat, mouse

**Positive Controls:** 

WB: HeLa cells, HepG2 cells, mouse kidney tissue, mouse liver tissue, mouse lung tissue, rat kidney tissue, NIH/3T3 cells, RAW 264.7 cells, HuH-7 cells, PC-

**Purification Method:** 

WB 1:2000-1:10000

Antigen affinity purification

Recommended Dilutions:

12 cells, C6 cells, human placenta tissue

## **Background Information**

Annexin A4 (Anxa4) is one of the Ca2 +-regulated and phospholipid-binding annexin superfamily proteins, and is located at 2p13. Anxa4 has a potential role in diagnosis, prognosis, and treatment of certain cancers. Annexin A4 shares 45 to 59% identity with other members and possibly interacts with ATP. Annexin A4 has in vitro anticoagulant activity and also inhibits phospholipase A2 activity. The Annexin A4 protein is expressed in a wide range of tissues, associated with polarized epithelial cells and is proposed to be involved in the regulation of chloride ions across the epithelial membrane.

Storage

Storage: Store at -20°C. Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29508-1-AP (Annexin A4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 29508-1-AP (Annexin IV antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.