For Research Use Only

Phospho-PERK/EIF2AK3 (Ser719) Polyclonal antibody



Catalog Number: 29546-1-AP

7 Publications

Basic Information

Catalog Number:

29546-1-AP

100ul , Concentration: 250 $\mu g/ml$ by

Nanodrop: Source:

Isotype:

Rabbit

IgG

GenBank Accession Number:

BC126354 GeneID (NCBI):

UNIPROT ID: Q9NZJ5

Full Name: eukaryotic translation initiation

factor 2-alpha kinase 3

Calculated MW: 1116 aa, 125 kDa Observed MW: 140 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

Species Specificity:

Human Cited Species: human, rat, mouse Positive Controls:

WB: λ phosphatase treated HeLa cells, Thapsigargin

treated HeLa cells

Background Information

EIF2AK3 encodes the protein kinase RNA-like ER kinase (PERK), a key regulator of the unfolded protein response (UPR) in response to ER stress. Under ER stress conditions, activation of PERK is triggered by the dissociation of glucose-regulated protein (GRP) 78 (also known as BiP) from its luminal domain, followed by oligomerization and autophosphorylation. Phosphorylated PERK subsequently phosphorylates eukaryotic translation initiation factor 2 alpha (eif2a), to attenuate global protein translation and reduce incoming ER protein load via upregulated ER chaperone expression. (PMID: 35922637, PMID: 32029570)

Notable Publications

Author	Pubmed ID	Journal	Application
Guanjun Li	38452464	Environ Int	WB
Yun Huang	38132921	Mar Drugs	WB
Jie Cui	37348685	Fish Shellfish Immunol	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Non-treated HeLa cells, phosphatase inhibitor treated and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 29546-1-AP (Phospho-PERI/EIF 2AK3 (Ser719) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Non-treated HeLa cells and Thapsigargin treated HeLa cells were subjected to SDS PAGE followed by western blot with 29546-1-AP (Phospho-PERK/EIF2AK3 (Ser719) antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.