For Research Use Only

BAK Polyclonal antibody

Catalog Number: 29552-1-AP

8 Publications



Basic Information

Catalog Number: GenBank Accession Number:

29552-1-AP NM 001188 GeneID (NCBI):

150ul , Concentration: 450 μ g/ml by 578

Nanodrop; Full Name:

Source: BCL2-antagonist/killer 1

Rabbit Calculated MW: Isotype: 23 kDa

IgG Observed MW: 23 kDa

Immunogen Catalog Number:

AG31042

Applications

Tested Applications:

IF, IHC, WB, ELISA **Cited Applications:**

IHC, WB

Species Specificity:

Human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A431 cells, Neuro-2a cells, HEK-293 cells, HepG2 cells, Jurkat cells, MOLT-4 cells, 4T1 cells, ROS1728

Purification Method:

WB 1:2000-1:12000 IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human colon cancer tissue,

IF: HeLa cells,

Background Information

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form oligomers or heterodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein localizes to mitochondria, and functions to induce apoptosis. It interacts with and accelerates the opening of the mitochondrial voltage-dependent anion channel, which leads to a loss in membrane potential and the release of cytochrome c. This protein also interacts with the tumor suppressor P53 after exposure to cell stress.

Notable Publications

Author	Pubmed ID	Journal	Application
Jingwen Tan	36208777	Chem Biol Interact	WB
Minghui Zhao	38238876	Animal Model Exp Med	WB
Ruimin Bai	37945816	Apoptosis	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29552-1-AP (BAK antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 29552-1-AP (BAK antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using BAK antibody (29552-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).