

For Research Use Only

# Phospho-MCL1 (Thr163) Polyclonal antibody



Catalog Number: 29560-1-AP

## Basic Information

<b>Catalog Number:</b> 29560-1-AP	<b>GenBank Accession Number:</b> BC017197	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 100ul , Concentration: 150 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 4170	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Source:</b> Rabbit	<b>Full Name:</b> myeloid cell leukemia sequence 1 (BCL2-related)	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 350 aa, 37 kDa	
	<b>Observed MW:</b> 40 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : MG132 treated HeLa cells,
<b>Species Specificity:</b> Human	

## Background Information

MCL1 is an anti-apoptotic member of the BCL-2 family originally isolated from the ML-1 human myeloid leukemia cell line. Similar to BCL2 and BCL2L1, MCL1 can interact with BAX and/or BAK1 to inhibit mitochondria-mediated apoptosis. Recent studies show that MCL1 is upregulated in numerous hematological and solid tumor malignancies. Therefore, MCL1 has been suggested as a potential new therapeutic target. MCL1 can be phosphorylated by several protein kinases which enables the recognition of MCL1 by its E3 ubiquitin-ligases TrCP or FBW7 (PMID: 33308268). MCL1 shows higher stability when phosphorylated on threonine 163 (PMID: 16543145).

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated and MG132 treated HeLa cells were subjected to SDS PAGE followed by western blot with 29560-1-AP (Phospho-MCL1 (Thr163) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as the loading control.