

For Research Use Only

Phospho-MCL1 (Thr163) Polyclonal antibody

Catalog Number: 29560-1-AP



Basic Information

Catalog Number:

29560-1-AP

Size:

100ul, Concentration: 150 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC017197

GeneID (NCBI):

4170

UNIPROT ID:

Q07820

Full Name:

myeloid cell leukemia sequence 1 (BCL2-related)

Calculated MW:

350 aa, 37 kDa

Observed MW:

40 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Positive Controls:

WB : MG132 treated HeLa cells,

Background Information

MCL1 is an anti-apoptotic member of the BCL-2 family originally isolated from the ML-1 human myeloid leukemia cell line. Similar to BCL2 and BCL2L1, MCL1 can interact with BAX and/or BAK1 to inhibit mitochondria-mediated apoptosis. Recent studies show that MCL1 is upregulated in numerous hematological and solid tumor malignancies. Therefore, MCL1 has been suggested as a potential new therapeutic target. MCL1 can be phosphorylated by several protein kinases which enables the recognition of MCL1 by its E3 ubiquitin-ligases TrCP or FBW7 (PMID: 33308268). MCL1 shows higher stability when phosphorylated on threonine 163 (PMID: 16543145).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and MG132 treated HeLa cells were subjected to SDS PAGE followed by western blot with 29560-1-AP (Phospho-MCL1 (Thr163) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as the loading control.