For Research Use Only

PDIR Polyclonal antibody

Catalog Number:29593-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number:

29593-1-AP BC001625 Antigen affinity purification
Size: GeneID (NCBI): Recommended Dilutions:

150ul , Concentration: 400 µg/ml by 10954 WB 1:2000-1:16000 Nanodrop; Full Name: IHC 1:50-1:500

Source: protein disulfide isomerase family A,

Rabbit member 5
Isotype: Calculated MW:
IgG 60 kDa
Immunogen Catalog Number: Observed MW:
AG31479 60 kDa

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IF, WB

Species Specificity: Human, Mouse, Rat Cited Species:

human

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, LO2 cells, HepG2 cells, mouse liver

Purification Method:

tissue, rat liver tissue

IHC: human placenta tissue,

Background Information

PDIR(Protein disulfide isomerase-related protein), also known as PDIA5 (Protein Disulfide Isomerase Family A Member 5), is a member of the PDI gene family and also exhibits chaperone-like activity. PDIR protein is correlated with immune infiltration, which is highly expressed in glioma and participates in glioma progression(PMID: 33664747). PDIR contributes to disulfide bond rearrangement in ATF6a under stress conditions that necessary for ATF6a activation upon ER stress(PMID: 24636989).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------|-----------|---------------|-------------|
| Yu Chen | 36003400 | Front Immunol | WB,IF |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29593-1-AP (PDIR antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 29593-1-AP (PDIR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 29593-1-AP (PDIR antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.