For Research Use Only

HTRA1 Polyclonal antibody

Catalog Number: 29618-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

29618-1-AP NM_002775 GeneID (NCBI): Size:

150ul , Concentration: 550 ug/ml by Nanodrop:

UNIPROT ID: Q92743 Rabbit Full Name:

Isotype: HtrA serine peptidase 1

IgG Calculated MW:

Immunogen Catalog Number: 51 kDa

AG31194 Observed MW:

51 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IHC 1:50-1:500 IF/ICC 1:200-1:800

Background Information

HTRA1, also named as HTRA, PRSS11 and L56, belongs to the peptidase S1B family. It is a protease that regulates the availability of nsulin-like growth factors (IGFs) by cleaving IGF-binding proteins. HTRA1 represses signaling by TGF-beta family members. Variations in the promoter region of HTRA1 are the cause of susceptibility to age-related macular degeneration type 7 (ARMD7). Defects in HTRA1 are the cause of cerebral autosomal recessive arteriopathy with subcortical infarcts and leukoencephalopathy (CARASIL). The 51 kDa HTRA1 was found at varying levels in the majority samples tested, whereas the 42 kDa form of HTRA1 was identified in degenerated intervertebral disc protein samples (PMID: 22556410). The short truncated monomer 37 kDa and dimer 74 kDa are proved (PMID:25329061). The antibody is specific to HTRA1.

Positive Controls:

IF/ICC: HepG2 cells,

WB: HeLa cells, human placenta tissue

IHC: human skin cancer tissue,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29618-1-AP (HTRA1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded skin cancer slide using 29618-1-AP (HTRA1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HTRA1 antibody (29618-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).