

For Research Use Only

DYNC2H1 Polyclonal antibody

Catalog Number: 29758-1-AP



Basic Information

Catalog Number: 29758-1-AP	GenBank Accession Number: NM_001377	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 ug/ml by Nanodrop;	GeneID (NCBI): 79659	Recommended Dilutions: WB 1:500-1:3000
Source: Rabbit	UNIPROT ID: Q8NCM8	IHC 1:50-1:500
Isotype: IgG	Full Name: dynein, cytoplasmic 2, heavy chain 1	IF/ICC 1:200-1:800
Immunogen Catalog Number: AG30707	Calculated MW: 493 kDa	
	Observed MW: 493 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: Human, mouse	WB : mouse testis tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse kidney tissue,
	IF/ICC : hTERT-RPE1 cells,

Background Information

DYNC2H1 (Cytoplasmic dynein 2 heavy chain 1) is involved in ciliary intraflagellar transport (IFT). DYNC2H1 drives retrograde transport of the IFT-A protein complex that regulates tip-to-base transport in cilia, involved in the generation and maintenance of cilia. DYNC2H1 variants or retina-predominant variants cause nonsyndromic retinal degeneration (PMID:32753734). DYNC2H1 mutations also cause asphyxiating thoracic dystrophy and short rib-polydactyly syndrome (PMID: 19442771). DYNC2H1 was localized to the basal bodies by immunofluorescence staining (PMID: 21552265, PMID: 26077881, PMID: 30320547).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 29758-1-AP (DYNC2H1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Methanol) fixed hTERT-RPE1 cells using DYNC2H1 antibody (29758-1-AP) at dilution of 1:400 and CoraLite® 594-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CoraLite® Plus 488 ARL13B antibody (CL488-17711, green).



Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using DYNC2H1 antibody (29758-1-AP) at dilution of 1:400 and CoraLite® 594-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CoraLite® Plus 488 ARL13B antibody (CL488-17711, green).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 29758-1-AP (DYNC2H1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 29758-1-AP (DYNC2H1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).