### For Research Use Only

# EBP50/NHERF-1 Polyclonal antibody

Catalog Number:29771-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

29771-1-AP NM 004252 GeneID (NCBI): Size:

150ul, Concentration: 600 ug/ml by

Nanodrop: **UNIPROT ID:** 014745 Rabbit

Full Name: Isotype: solute carrier family 9 (sodium/hydrogen exchanger), IgG

member 3 regulator 1 Immunogen Catalog Number:

AG30878 Calculated MW:

39 kDa Observed MW: 50 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:1000-1:4000 IF/ICC 1:200-1:800

### **Applications**

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

WB: HCT 116 cells, HepG2 cells, MCF-7 cells, MDA-MB-231 cells, rat kidney tissue, mouse kidney tissue IHC: human placenta tissue, rat kidney tissue

IF/ICC: HeLa cells, A549 cells

## **Background Information**

Na+/H+ exchanger regulatory factor 1 (NHERF-1, also known as EBP50 ezrin-binding protein of 50 kDa, SLCA9A3R1) is a phosphoprotein that assembles multiprotein complexes via two PDZ domains and a C-terminal ezrin-binding domain. NHERF-1 was initially recognized as a scaffolding protein that recruits membrane transporters/receptors and cytoplasmic signaling proteins into functional complexes localized at or near the plasma membrane in epithelial cells (PMID: 19308292).

### Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 29771-1-AP (EBP50/NHERF-1 antibody) at dilution of 1:10000 and incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 29771-1-AP (EBP50/NHERF-1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 29771-1-AP (EBP50/NHERF-1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 29771-1-AP (EBP50/NHERF-1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 29771-1-AP (EBP50/NHERF-1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using EBP50/NHERF-1 antibody (29771-1-AP) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using EBP50/NHERF-1 antibody (29771-1-AP) at dilution of 1:600 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-HUMAN EBP50/NHERF-1 (29771-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).