For Research Use Only

PD-1/CD279 Polyclonal antibody Catalog Number:29780-1-AP




Basic Information	Catalog Number: 29780-1-AP	GenBank Accession Number: BC074740	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 450 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG31288	GeneID (NCBI): 5133 UNIPROT ID: Q15116 Full Name: programmed cell death 1 Calculated MW: 288 aa, 32 kDa Observed MW: 55 kDa	Recommended Dilutions: WB 1:500-1:3000
Applications	Tested Applications: WB, ELISA Species Specificity: Human, Mouse, Rat	Positive Controls: WB : mouse thymus tissue, rat thymus tissue	
Background Information	Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436). It has been reported that PD-1 is heavily glycosylated and migrates with an apparent molecular mass of 47-55 kDa on SDS-PAGE , which is larger than its predicted mass of 32 kDa (PMID: 8671665; 17640856; 17003438).		
Storage	Storage: Store at -20°C. Storage Buffer: PBS with 0.02% sodium azide and 50	0, 1	
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29780-1-AP (PD-1/CD279 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.