## For Research Use Only

## MGC33407/ACTL9 Polyclonal antibody

Catalog Number: 29885-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

29885-1-AP BC024028 GeneID (NCBI): Size: 150ul , Concentration: 450 ug/ml by 284382 Nanodrop: **UNIPROT ID:** 

Q8TC94 Rabbit Full Name:

Isotype: hypothetical protein MGC33407

IgG Observed MW: Immunogen Catalog Number: 45 kDa

AG31805

**Applications** 

**Tested Applications:** 

WB, IHC, IF-P, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, rat skeletal muscle

**Purification Method:** 

WB 1:500-1:2000

IHC 1:50-1:500

IF-P 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human liver tissue, mouse testis tissue

IF-P: mouse testis tissue,

## **Background Information**

ACTL9 (Actin Like 9), also named MGC33407, encodes a testis-specific actin-like protein that plays a role in fusion of proacrosomal vesicles and perinuclear theca formation (PMID: 33626338). ACTL9 protein contains 416 amino acids, which mainly localizes in the equatorial segment of human sperm head (PubMed: 33626338). Diseases associated with ACTL9 include spermatogenic failure and non-syndromic male infertility.

Storage


Storage: Store at -20°C. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

\*\*\* 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 29885-1-AP (MGC33407/ACTL9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 29885-1-AP (MGC33407/ACTL9 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 29885-1-AP (MGC33407/ACTL9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using MGC33407/ACTL9 antibody (29885-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse testis tissue using MGC33407/ACTL9 antibody (29885-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 29885-1-AP (MGC 33407/ACTL9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 29885-1-AP (MGC33407/ACTL9 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).