## For Research Use Only

## BAHCC1 Polyclonal antibody

Catalog Number: 30046-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

30046-1-AP NM\_001291324.1 Size: GeneID (NCBI):

150ul , Concentration: 500 ug/ml by 57597
Nanodrop; UNIPROT ID:

Source: Q9P281
Rabbit Full Name:

Isotype: BAH domain and coiled-coil

IgG containing 1
Immunogen Catalog Number: Calculated MW:
AG32599 280 kDa
Observed MW:

280 kDa

**Applications** 

Tested Applications:

IHC, IF/ICC, ELISA

Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

IHC: mouse brain tissue, rat brain tissue, mouse colon

**Purification Method:** 

IHC 1:50-1:500 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

tissue

IF/ICC: A549 cells,

## **Background Information**

BAHCC1 (BAH and coiled-coil domain-containing protein 1) is also named as BAHD2, KIAA1447 and BAH domain-containing protein 2. BAHCC1 is a transcriptional regulator controlling expression of E2F/KLF-dependent cell-cycle and DNA-repair genes. BAHCC1 associates with BRG1-containing remodeling complexes at the promoters of these genes. BAHCC1 silencing leads to decreased cell proliferation and delayed DNA repair. Consequently, BAHCC1 deficiency cooperates with PARP inhibition to induce melanoma cell death (PMID: 37924516). depletion of BAHCC1, or disruption of the BAHCC1BAH-H3K27me3 interaction, causes derepression of H3K27me3-targeted genes that are involved in tumor suppression and cell differentiation, leading to suppression of oncogenesis (PMID: 33139953).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A549 cells using BAHCC1 antibody (30046-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 30046-1-AP (BAHCC1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).