

For Research Use Only

POGZ Polyclonal antibody

Catalog Number: 30106-1-AP



Basic Information

Catalog Number: 30106-1-AP	GenBank Accession Number: NM_015100	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop;	GeneID (NCBI): 23126	Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for IP and 1:1000-1:4000 for WB
Source: Rabbit	Full Name: pogo transposable element with ZNF domain	IHC 1:50-1:500 IF 1:200-1:800
Isotype: IgG	Calculated MW: 155kd	
Immunogen Catalog Number: AG32435	Observed MW: 155 kDa	

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Species Specificity:

Human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HepG2 cells, SH-SY5Y cells, mouse brain tissue

IP : HeLa cells,

IHC : mouse brain tissue, mouse testis tissue

IF : HepG2 cells, U-251 cells

Background Information

POGZ, also known as MRD37, is a zinc-finger protein containing 9 C2H2 zinc fingers, a CENP-B (centromere protein-B) domain, and a DDE domain found in the bacterial transposase/retroviral integrase family. POGZ plays a role in mitotic cell cycle progression and is involved in kinetochore assembly and mitotic sister chromatid cohesion. POGZ plays a role in mitotic chromosome segregation by regulating aurora kinase B/AURKB activation and AURKB and CBX5 dissociation from chromosome arms (PMID:20562864). Heterozygous mutation of POGZ causes White-Sutton syndrome (PMID: 26739615). Expect a band approximately 155 kDa in size corresponding to POGZ by western blotting.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30106-1-AP (POGZ antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using POGZ antibody (30106-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




IP result of anti-POGZ(IP:30106-1-AP, 4ug; Detection:30106-1-AP 1:2000) with HeLa cells lysate 1800 ug.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 30106-1-AP (POGZ antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using POGZ antibody (30106-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).