

For Research Use Only

# NFATC1 Polyclonal antibody

Catalog Number: 30114-1-AP



## Basic Information

<b>Catalog Number:</b> 30114-1-AP	<b>GenBank Accession Number:</b> BC104753	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 650 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 4772	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IF/ICC 1:200-1:800
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> O95644	
<b>Isotype:</b> IgG	<b>Full Name:</b> nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1	
<b>Immunogen Catalog Number:</b> AG32365	<b>Calculated MW:</b> 716aa,78 kDa; 943aa,101 kDa	
	<b>Observed MW:</b> 82-140 kDa	

## Applications

<b>Tested Applications:</b> WB, IF/ICC, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> Human	<b>WB :</b> HEK-293 cells, <b>IF/ICC :</b> U-251 cells,

## Background Information

Nuclear factor of activated T-cells cytoplasmic (NFATC) is a family of transcription factors originally identified in T-cells. The family has five members (NFATC1 through NFATC5), which play roles both within and outside the immune system. NFATC1 is widely expressed and resides in endocardial cushion endothelial cells, skeletal muscle fibers, T lymphocytes, osteoblasts, osteoclasts, etc. The autoregulation of NFATC1 is a crucial step for cell fate determination. It is a master regulator of RANKL-induced osteoclast differentiation and plays a pivotal role in osteoclast fusion and osteoclast activation via up-regulation of various genes responsible for osteoclast adhesion, migration, acidification, degradation of inorganic and organic bone matrix (PMID: 20035895, PMID: 19754901). NFAT proteins is alternatively modified to generate splice variants. NFATC1 can be detected isoforms of 82-140 kDa (PMID: 18097033).

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30114-1-AP (NFATC1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using NFATC1 antibody (30114-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).