For Research Use Only

BST2 Polyclonal antibody

Catalog Number:30118-1-AP

Featured Product




Basic Information	Catalog Number: 30118-1-AP	GenBank Accession Number: BC033873	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG32333	GeneID (NCBI): 684 UNIPROT ID: Q10589 Full Name: bone marrow stromal cell antiger Calculated MW: 180 aa, 20 kDa Observed MW: 30-36 kDa, 50-70 kDa	Recommended Dilutions: WB 1:5000-1:50000 IF-P 1:200-1:800
Applications	Tested Applications: WB, IF-P, ELISA Species Specificity: Human, mouse	Positive C WB : HeLa IF-P : mou	
Background Information	BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).		
Storage *** 20ul sizes contain 0.1% BSA	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50 Aliquoting is unnecessary for -20°C s	% glycerol pH 7.3.	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 30118-1-AP (BST2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of BST2 antibody (30118-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-BST2 transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using BST2 antibody (30118-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using BST2 antibody (30118-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).