For Research Use Only

BRAWNIN Polyclonal antibody

Catalog Number:30212-1-AP



Purification Method:

WB 1:400-1:1000

IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

30212-1-AP GeneID (NCBI): Size: 728568 150ul, Concentration: 400 ug/ml by **UNIPROT ID:** Nanodrop:

Q69YU5 Full Name:

Rabbit chromosome 12 open reading frame

Isotype:

IgG Calculated MW:

Immunogen Catalog Number: 8 kDa

AG32955 Observed MW:

9 kDa

Applications

Positive Controls: **Tested Applications:**

WB, IF/ICC, ELISA WB: K-562 cells, MCF-7 cells Species Specificity: IF/ICC: U-251 cells, A431 cells Human

Background Information

Ubiquinol-cytochrome-c reductase complex assembly factor 6 (UQCC6) is also named as BRAWNIN, BR and C12orf73, and belongs to the UQCC6 family. BRAWNIN (BR), a 71 a.a. peptide encoded by C12orf73, is essential for respiratory chain complex III (CIII) assembly. In human cells, BRAWNIN is induced by the energy-sensing AMPK pathway, and its depletion impairs mitochondrial ATP production. In zebrafish, Brawnin deletion causes complete CIII loss, resulting in severe growth retardation, lactic acidosis and early death (PMID:32161263). BR is also detectable in human cardiac and skeletal muscle where it displays a staining pattern characteristic of the mitochondria network. BR protein abundance in mouse tissues correlated well with that of mitochondrial respiratory chain proteins, being high in brown adipose, cardiac and skeletal muscle but virtually undetectable in white $adipose\ tissue\ (PMID: 32161263).\ BR\ resides\ in\ the\ IMM\ and\ not\ in\ the\ outer\ mitochondrial\ membrane\ (OMM)$ (PMID:32161263).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30212-1-AP (BRAWNIN antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using BRAWNIN antibody (30212-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using BRAWNIN antibody (30212-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed A431 cells using BRAWNIN antibody (30212-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).