## For Research Use Only

## Phospho-EGFR (Tyr1069) Polyclonal antibody

Catalog Number: 30277-1-AP

4 Publications




Basic Information	Catalog Number: 30277-1-AP	GenBank Accession Number: BC094761	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	100ul , Concentration: 350 ug/ml by	1956	WB 1:500-1:2000	
	Nanodrop; Source:	UNIPROT ID: P00533		
	Rabbit	Full Name:		
	Isotype: IgG	epidermal growth factor receptor (erythroblastic leukemia viral (v-erb- b) oncogene homolog, avian)		
		Calculated MW: 1210 aa, 134 kDa		
		Observed MW: 170 kDa		
Applications	Tested Applications: WB, ELISA	Positive Controls:		
	Cited Applications: WB, IF	WB : EGF treated HeLa cells,		
	Species Specificity: Human			
	Cited Species:			
	human			
Background Information	human Epidermal Growth Factor Receptor (E family which consists of ErbB1 (EGFR activation of EGFR is among the mos largely through different classes of g acts as the degron of EGFR and is reco	t or HER1), ErbB2 (HER2 or Neu), ErbB3 t common oncogenic driving events i enomic alterations within the EGFR g ognized by c-Cbl, enhancing the ubiq F stimulation. (PMID: 35164092, PMI	(HER3), and ErbB4 (HER4). Aberrant n human cancer, which is mediated gene. Phosphorylation of EGFR Tyr1069 uitination and degradation of EGFR. EGF D: 29464683, PMID: 37085908). Some	
	human Epidermal Growth Factor Receptor (E family which consists of ErbB1 (EGFR activation of EGFR is among the mos largely through different classes of g acts as the degron of EGFR and is rece stability is tightly regulated upon EG scientists call EGFR Tyr1069 as EGFR	t or HER1), ErbB2 (HER2 or Neu), ErbB3 t common oncogenic driving events i enomic alterations within the EGFR g ognized by c-Cbl, enhancing the ubiq F stimulation. (PMID: 35164092, PMI	n human cancer, which is mediated gene. Phosphorylation of EGFR Tyr1069 uitination and degradation of EGFR. EGFI D: 29464683, PMID: 37085908). Some	
	human Epidermal Growth Factor Receptor (E family which consists of ErbB1 (EGFR activation of EGFR is among the mos largely through different classes of g acts as the degron of EGFR and is rec stability is tightly regulated upon EG scientists call EGFR Tyr1069 as EGFR Author	t or HER1), ErbB2 (HER2 or Neu), ErbB3 t common oncogenic driving events i enomic alterations within the EGFR g ognized by c-Cbl, enhancing the ubiq IF stimulation. (PMID: 35164092, PMI Tyr1045, the site where the signaling	(HER3), and ErbB4 (HER4). Aberrant n human cancer, which is mediated gene. Phosphorylation of EGFR Tyr1069 uitination and degradation of EGFR. EGF D: 29464683, PMID: 37085908). Some g peptide is removed.	
	human Epidermal Growth Factor Receptor (E family which consists of ErbB1 (EGFR activation of EGFR is among the mos largely through different classes of g acts as the degron of EGFR and is rece stability is tightly regulated upon EG scientists call EGFR Tyr1069 as EGFR Author Pub Jie Yuan 398	t or HER1), ErbB2 (HER2 or Neu), ErbB3 t common oncogenic driving events i enomic alterations within the EGFR g ognized by c-Cbl, enhancing the ubiq F stimulation. (PMID: 35164092, PMI Tyr1045, the site where the signaling med ID Journal	(HER3), and ErbB4 (HER4). Aberrant n human cancer, which is mediated gene. Phosphorylation of EGFR Tyr1069 uitination and degradation of EGFR. EGF D: 29464683, PMID: 37085908). Some g peptide is removed. Application	
Background Information Notable Publications	human Epidermal Growth Factor Receptor (E family which consists of ErbB1 (EGFR activation of EGFR is among the mos largely through different classes of g acts as the degron of EGFR and is rece stability is tightly regulated upon EC scientists call EGFR Tyr1069 as EGFR Author Put Jie Yuan 398 Bingling Zhang 397	tor HER1), ErbB2 (HER2 or Neu), ErbB3 t common oncogenic driving events i enomic alterations within the EGFR g ognized by c-Cbl, enhancing the ubiqu F stimulation. (PMID: 35164092, PMI Tyr1045, the site where the signaling med ID Journal 17673 IUBMB Life	(HER3), and ErbB4 (HER4). Aberrant n human cancer, which is mediated gene. Phosphorylation of EGFR Tyr1069 uitination and degradation of EGFR. EGF D: 29464683, PMID: 37085908). Some g peptide is removed. Application WB	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated HeLa and EGF treated HeLa cells were subjected to SDS PAGE followed by western blot with 30277-1-AP (Phospho-EGFR (Tyr1069) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Vinculin antibody as loading control.