

For Research Use Only

IFN Gamma Polyclonal antibody

Catalog Number: 30293-1-AP



Basic Information

Catalog Number: 30293-1-AP	GenBank Accession Number: NM_138880	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop;	GeneID (NCBI): 25712	Recommended Dilutions: IHC 1:50-1:500
Source: Rabbit	Full Name: interferon gamma	
Isotype: IgG	Calculated MW: 18 kDa	

Applications

Tested Applications: FC, IHC, ELISA	Positive Controls: IHC : rat spleen tissue,
Species Specificity: rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IFN gamma (IFNG) is a soluble cytokine that is the only member of the type II class of IFN. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has been identified in many studies as a biomarker for pleural tuberculosis (TB).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1X10⁶ PMA and ionomycin treated rat splenocytes cells were intracellularly stained with 0.5 ug Anti-Rat IFN Gamma (30293-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000, and 0.5 ug APC Anti-Rat CD4 (OX-35) (APC-65179, Clone: OX-35), and 0.5 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm



Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using 30293-1-AP (IFN Gamma antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using 30293-1-AP (IFN Gamma antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).