#### For Research Use Only

# TRIM46 Polyclonal antibody

Catalog Number: 30298-1-AP 1 Publications



**Basic Information** 

Catalog Number: 30298-1-AP

Size:

GenBank Accession Number:

BC069416

GeneID (NCBI):

80128

150ul, Concentration: 800 ug/ml by Nanodrop:

**UNIPROT ID:** Q7Z4K8

Rabbit Full Name:

Isotype: tripartite motif-containing 46

IgG Calculated MW: Immunogen Catalog Number: 759 aa, 83 kDa AG33175 Observed MW:

83 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:500-1:3000

IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** WB, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity: mouse, rat, human **Cited Species:** mouse

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IF/ICC: HepG2 cells,

## **Background Information**

Tripartite motif-containing protein 46 (TRIM46), a microtubule-associated protein, is involved in the formation of parallel microtubule bundles linked by cross-bridges in the proximal axon. Required for the uniform orientation and maintenance of the parallel microtubule fascicles, which are important for efficient cargo delivery and trafficking in axons. Thereby also required for proper axon specification, the establishment of neuronal polarity, and proper neuronal migration. The TRIM46 protein reported in previous studies includes the long isoform TRIM46L (83 kDa) and the short isoform TRIM46S (60 kDa).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Anja Konietzny	38525600	J Cell Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

### Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30298-1-AP (TRIM46 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using TRIM46 antibody (30298-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).