

For Research Use Only

OGG1 Polyclonal antibody

Catalog Number: 30302-1-AP



Basic Information

Catalog Number: 30302-1-AP	GenBank Accession Number: BC000657	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 4968	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	Full Name: 8-oxoguanine DNA glycosylase	
Isotype: IgG	Calculated MW: 22 kDa, 36-40 kDa, 45-57 kDa	
Immunogen Catalog Number: AG32962	Observed MW: 36-40 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, rat skeletal muscle tissue
Species Specificity: Human, Rat	

Background Information

The DNA damages induced by ROS contain base modification, base loss, and DNA single strand breaks, which are usually repaired by the base excision repair (BER) pathway in both prokaryotes and eukaryotes. OGG1 (The human 8-oxoguanine glycosylase 1) is the primary enzyme in BER pathway, responsible for the excision of 7, 8-dihydro-8-oxoguanine (8-oxoG), a mutagenic base byproduct that occurs as a result of exposure to reactive oxygen species. There's 8 isoforms of OGG1, with calculated MW 22 kDa, 36-40 kDa and 45-57 kDa. The difference among these isoforms is the C-terminal (317-345aa).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30302-1-AP (OGG1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.