For Research Use Only

CDX2 Polyclonal antibody

Catalog Number: 30380-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

30380-1-AP BC014461 GeneID (NCBI): Size:

Nanodrop: **UNIPROT ID:** Q99626 Rabbit Full Name:

150ul , Concentration: 600 µg/ml by 1045

Isotype: caudal type homeobox 2

IgG Calculated MW: Immunogen Catalog Number: 313 aa, 34 kDa AG32826 Observed MW: 33-40 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500 IF 1:50-1:500

Applications

Tested Applications: WB, IF, IHC, ELISA

Species Specificity:

Human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse colon tissue, pig colon tissue

IHC: human colon cancer tissue, mouse colon tissue,

rat colon tissue IF: Caco-2 cells,

Background Information

CDX2, also named as Homeobox protein CDX-2, is a 313 amino acid protein, which contains one homeobox DNAbinding domain and belongs to the Caudal homeobox family. CDX2 localizes in the nucleus and is involved in the transcriptional regulation of multiple genes expression in the intestinal epithelium. The relative expression of CDX1 to CDX2 protein may be important in the anterior to posterior patterning of the intestinal epithelium and in defining patterns of proliferation and differentiation along the crypt-villus axis. Both Cdx1 and Cdx2 genes must be expressed to reduce tumorigenic potential, to increase sensitivity to apoptosis, and to reduce cell migration, suggesting that the two genes control the normal phenotype by independent pathways.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30380-1-AP (CDX2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 30380-1-AP (CDX2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 30380-1-AP (CDX2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 30380-1-AP (CDX2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 30380-1-AP (CDX2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using CDX2 antibody (30380-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 30380-1-AP (CDX2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).