For Research Use Only

SYNCRIP Polyclonal antibody

Catalog Number: 30443-1-AP



Purification Method:

WB 1:1000-1:8000 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

30443-1-AP BC032643 GeneID (NCBI): Size: 150ul, Concentration: 500 ug/ml by 10492

Nanodrop: **UNIPROT ID:** 060506 Rabbit Full Name:

Isotype: synaptotagmin binding, cytoplasmic

RNA interacting protein IgG

Immunogen Catalog Number: Calculated MW: 623 aa, 70 kDa AG31509

> Observed MW: 62-74 kDa

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA WB: K-562 cells, THP-1 cells Species Specificity: IF/ICC: A431 cells,

Background Information

 $Synaptotagmin-binding, cytoplasmic\ RNA-interacting\ protein (SYNCRIP), also\ identified\ as\ hnRNP\ Q,\ involves\ in$ mRNA processing mechanisms. It's component of the CRD-mediated complex that promotes MYC mRNA stability. Also it is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major codingregion determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. As previously reported, the 65 and 74 kDa proteins corresponded to two isoforms of SYNCRIP produced from alternatively spliced mRNAs (PMID: 15340051).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30443-1-AP (SYNCRIP antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using SYNCRIP antibody (30443-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).