For Research Use Only

## NUCB2 Polyclonal antibody Catalog Number: 30604-1-AP




Basic Information	Catalog Number: 30604-1-AP	GenBank Accession Number: NM_005013	Purification Method: Antigen affinity purification		
	Size: 150ul , Concentration: 800 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG30704	GeneID (NCBI): 4925 UNIPROT ID: P80303 Full Name: nucleobindin 2	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:200-1:800 IF/ICC 1:200-1:800		
				Calculated MW: 50 kDa	
				Observed MW: 46-50 kDa	
		Applications			ve Controls:
					use brain tissue, human milk, SGC-7901 cells, ı tissue
human, mouse, rat Note-IHC: suggested antigen 1			uman intrahepatic cholangiocarcinoma tissue,		
TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			: U2OS cells,		
Background Information	NucB2, an anorexigenic molecule, is expressed mainly in the hypothalamus, particularly in the supraoptic nucleus (SON) and the paraventricular nucleus (PVN). NucB2 is also expressed in the subfornical organ (SFO). Because the SON and PVN are involved in body fluid regulation, NucB2 may be involved in dehydration-induced anorexia. NUCB2 is composed of a signal peptide of 24 amino acids in the N-terminal region and a protin structure containing 396 amino acids. Nesfatin-1 is an 82-amino-acid peptide hormone cleaved from nucleobindin2 (NUCB2).				
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer:				
	PBS with 0.02% sodium azide and 50 Aliquoting is unnecessary for $-20^{\circ}$ C s				

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 30604-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 30604-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-Immunonstochemical analysis of paramite embedded human intrahepatic cholangiocarcinoma tissue slide using 30604-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using NUCB2/nesfatin-1 antibody (30604-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).