

For Research Use Only

EPOR Polyclonal antibody

Catalog Number: 30766-1-AP



Basic Information

Catalog Number: 30766-1-AP	GenBank Accession Number: NM_000121.4	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 450 µg/ml by Nanodrop;	GeneID (NCBI): 2057	Recommended Dilutions: WB 1:1000-1:5000
Source: Rabbit	Full Name: erythropoietin receptor	
Isotype: IgG	Calculated MW: 55 kDa	
Immunogen Catalog Number: AG33956	Observed MW: 70 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human placenta tissue, mouse kidney tissue, mouse spleen tissue
Species Specificity: Human, Mouse	

Background Information

Erythropoietin (EPO) receptor (EPOR) is a glycoprotein that belongs to the type I superfamily of single-transmembrane cytokine receptors. It consists of an extracellular domain that binds to the EPO ligand, a transmembrane domain and an intracellular domain. The interaction of EPO and EPOR triggers the activation of several signaling pathways that induce erythropoiesis, including JAK2/STAT5, PI3K/AKT, and MAPK (PMID: 34281163). EPOR is present on erythroid progenitor cells and has also been detected on a wide variety of non-hematopoietic cells (PMID: 36233351). The calculated molecular weight of EPOR is 55 kDa. EPOR undergoes posttranslational modification and the apparent molecular weight of 66-78 kDa has been reported (PMID: 8943308; 12088257).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30766-1-AP (EPOR antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.