For Research Use Only

CD14 Polyclonal antibody

Catalog Number: 30796-1-AP



Purification Method:

WB 1:500-1:1000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

GenBank Accession Number:

BC010507

30796-1-AP Size:

Nanodrop;

GeneID (NCBI):

150ul, Concentration: 500 ug/ml by

ENSEMBL Gene ID: ENSG00000170458

Source: Rabbit

UNIPROT ID: P08571

Isotype: IgG

Full Name: CD14 molecule

Calculated MW: 375 aa, 40 kDa

Observed MW:

50-55 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HL-60 cells,

IHC: human tonsillitis tissue,

Background Information

CD14 is a 50-55 kDa glycosylphosphatidylinositol-anchored glycoprotein preferentially expressed on monocytes and macrophages, and at lower levels on granulocytes (PMID: 3385210; 2462937; 7685797). CD14 can also exist as a soluble protein. CD14 acts as a co-receptor for bacterial liposaccharides (LPS) (PMID: 1698311). It plays a major role in the inflammatory response of monocytes to LPS.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



HL-60 cells were subjected to SDS PAGE followed by western blot with 30796-1-AP (CD14 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 30796-1-AP (CD14 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).