For Research Use Only

CD68 Polyclonal antibody

Catalog Number: 30929-1-AP 1 Publications



Basic Information

Catalog Number:

30929-1-AP NM_001291058.1

GeneID (NCBI): Size:

150ul, Concentration: 500 ug/ml by 12514 Nanodrop;

Source: Rabbit Isotype: IgG

Calculated MW:

35 kDa

UNIPROT ID:

P31996-1

Full Name:

CD68 antigen

GenBank Accession Number:

Antigen affinity purification Recommended Dilutions: WB 1:2000-1:12000

Purification Method:

IHC 1:300-1:1200 IF-P 1:50-1:500

Applications

Tested Applications: WB, IHC, IF-P, ELISA **Cited Applications:**

IHC

Species Specificity: mouse, rat **Cited Species:** mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: NR8383 cells, RAW 264.7 cells

IHC: mouse spleen tissue, mouse liver tissue

IF-P: mouse spleen tissue,

Background Information

Mouse CD68 (also known as macrosialin) is a type I transmembrane glycoprotein that is highly and specifically expressed by mouse tissue macrophages, and to a lesser extent by dendritic cells. It belongs to the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. CD68 is also a member of the scavenger receptor family. It may play a role in phagocytic activities of tissue macrophages.

Notable Publications

Author	Pubmed ID	Journal	Application
Yafei Tian	39609656	J Nanobiotechnology	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30929-1-AP (CD68 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 30929-1-AP (CD68 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 30929-1-AP (CD68 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using CD68 antibody (30929-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using CD68 antibody (30929-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).