

For Research Use Only

IL-12/IL-23 p40 Polyclonal antibody

Catalog Number: 30956-1-AP



Basic Information

Catalog Number: 30956-1-AP	GenBank Accession Number: BC067498	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop;	GeneID (NCBI): 3593	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	Full Name: interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40)	
Isotype: IgG	Calculated MW: 328 aa, 37 kDa	
	Observed MW: 40-45 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Recombinant protein,
Species Specificity: Human	

Background Information

IL-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R β 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Human IL-12B/p40, IL-12/p70, IL-12A/p35 recombinant protein, were subjected to SDS PAGE followed by western blot with 30956-1-AP (IL-12B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.