

For Research Use Only

NRG1, isoform Alpha Polyclonal antibody



Catalog Number: 31043-1-AP

Basic Information

Catalog Number: 31043-1-AP	GenBank Accession Number: BC007675	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 800 µg/ml by Nanodrop;	GeneID (NCBI): 3084	Recommended Dilutions: WB 1:500-1:1000 IF 1:200-1:800
Source: Rabbit	Full Name: neuregulin 1	
Isotype: IgG	Calculated MW: 32 kDa	
Immunogen Catalog Number: AG35114	Observed MW: 70 kDa	

Applications

Tested Applications:
IF, WB, ELISA

Species Specificity:
Human, Mouse, Rat

Positive Controls:

WB: A431 cells, A549 cells, HeLa cells, MCF-7 cells, SH-SY5Y cells, mouse brain tissue, rat brain tissue

IF: MCF-7 cells,

Background Information

Neuregulin 1 (NRG1) is a trophic factor that has been implicated in neural development, neurotransmission, and synaptic plasticity. NRG1 has multiple isoforms that are generated by usage of different promoters and alternative splicing of a single gene. NRG1 and its receptor ErbB tyrosine kinase are expressed not only in the developing nervous system, but also in the adult brain. The antibody is specific to NRG1 isoform Alpha.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using NRG1, isoform Alpha antibody (31043-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 31043-1-AP (NRG1, isoform Alpha antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.