For Research Use Only

EGR1 Polyclonal antibody

Catalog Number: 31098-1-AP



Purification Method:

WB 1:500-1:2000

Antigen affinity Purification

Recommended Dilutions:

Basic Information

Catalog Number:

GenBank Accession Number:

BC073983

31098-1-AP Size:

GeneID (NCBI):

Genera (NCBI):

150ul, Concentration: 500 ug/ml by 1958

1958

Nanodrop;

UNIPROT ID: P18146

Rabbit

Full Name:

Isotype: IgG early growth response 1
Calculated MW:

Immunogen Catalog Number:

543 aa, 58 kDa

AG33440

Observed MW:

70-80 kDa

Applications

Tested Applications:

WB, ELISA

Positive Controls:

WB: A549 cells, HeLa cells, Jurkat cells

Species Specificity: Human

Background Information

EGR1, also named as Early growth response protein 1, is a 543 amino acid protein, which belongs to the EGR C2H2-type zinc-finger protein family. EGR1 as a transcriptional regulator recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3'(EGR-site) in the promoter region of target genes and binds double-stranded target DNA, irrespective of the cytosine methylation status. EGR1 regulates the transcription of numerous target genes, and thereby plays an important role in regulating the response to growth factors, DNA damage, and ischemia. EGR1 activates expression of p53/TP53 and TGFB1, and thereby helps prevent tumor formation and plays a role in the regulation of cell survival, proliferation and cell death. Egr1 is suggested to be a 55-kDa protein according to the translation of its coding sequence. Analysis of cytoplasmic and nuclear fractions of normal and irradiated native CNS tissue by Western blotting revealed a 110-kDa band for Egr1 localized in the cytoplasm and a 75-kDa version for the nuclear Egr1(PMID: 17497096)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31098-1-AP (EGR1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.