## For Research Use Only Cullin 5 Polyclonal antibody

Catalog Number:31183-1-AP




Basic Information	Catalog Number: 31183-1-AP	GenBank Accession Number: BC063306	Purification Method: Antigen affinity Purification
	Size: 150ul, Concentration: 250 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG34620	GeneID (NCBI): 8065 UNIPROT ID: Q93034 Full Name: cullin 5 Calculated MW: 91 kDa Observed MW: 80 kDa, 90 kDa	Recommended Dilutions: WB 1:1000-1:10000 IHC 1:50-1:500
Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	ively, antigen	
Background Information	The cullin family proteins are scaffold proteins for the RING finger type E3 ligases, participating in the proteolysis through the ubiquitin-proteasome pathway. In humans, there are eight members of cullins, including CUL1, 2, 3, 4A, 4B, 5, 7, and CUL9 with an evolutionarily conserved cullin homology domain at the C-terminus, responsible for their interaction with the RING protein, assembling 8 cullin-based RING dependent E3 ubiquitin ligases to target thousands of substrates for ubiquitination and degradation (PMID: 32334051). Cullin 5 is the key scaffolding protein of Cullin RING ligase-5 (CRL-5). Cullin 5 plays positive roles in regulating cell growth, proliferation and physiological and other processes in the human body (PMID: 32194910). Cullin 5 has been detected in the form of double bands in the results of Western blot, which might be caused by neddylation (PMID: 36449095; 31294695).		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	% glycerol pH 7.3.	
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^\circ$ C s	itorage	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 31183-1-AP (Cullin 5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 31183-1-AP (Cullin 5 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31183-1-AP (Cullin 5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31183-1-AP (Cullin 5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).