## For Research Use Only

## Apoe Polyclonal antibody

Catalog Number: 31454-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number: 31454-1-AP NM\_001305819.1

GeneID (NCBI): Size:

150ul, Concentration: 300 ug/ml by 11816 Nanodrop: UNIPROT ID:

P08226 Rabbit Full Name: Isotype: Apolipoprotein E IgG Calculated MW:

Immunogen Catalog Number: 36kDa EG1093

Observed MW:

31~36 kDa

**Purification Method:** 

Antigen affinity Purification Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:300-1:1200 IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA Species Specificity:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: U-251 cells, BV-2 cells, C2C12 cells, mouse brain tissue, mouse skeletal muscle tissue, mouse cerebellum tissue, rat brain tissue, pig cerebellum

IHC: rat cerebellum tissue, IF/ICC: HepG2 cells,

## **Background Information**

Apolipoprotein E (Apoe) is a core component of plasma lipoproteins that participates in diverse biological processes, including plasma lipoprotein metabolism, intracellular cholesterol utilization, cell growth, immunoregulation, and neuronal growth and repair (PMID: 17854398). Mouse apoE, like apoE4, contains the equivalent of Arg-112 and Glu-255, but lacks the critical Arg-61 equivalent (it contains Thr-61). Thus, mouse apoE does not display apoE4 domain interaction (PMID: 11553788).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 314,54-1-AP (Apoe antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 31454-1-AP (Apoe antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded rat cerebellum tissue slide using 31454-1-AP (Apoe antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat cerebellum tissue slide using 31454-1-AP (Apoe antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Apoe antibody (31454-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).