

For Research Use Only

TFPI Polyclonal antibody

Catalog Number: 31765-1-AP



Basic Information

Catalog Number: 31765-1-AP	GenBank Accession Number: NM_006287.6	Purification Method: Antigen affinity Purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop;	GeneID (NCBI): 7035	Recommended Dilutions: WB 1:500-1:3000
Source: Rabbit	UNIPROT ID: P10646-1	
Isotype: IgG	Full Name: tissue factor pathway inhibitor (lipoprotein-associated coagulation inhibitor)	
	Calculated MW: 35 kDa	
	Observed MW: 35-40 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HepG2 cells,
Species Specificity: human	

Background Information

Tissue factor pathway inhibitor (TFPI) is an anticoagulant protein produced primarily by the endothelium and megakaryocytes and alternative mRNA splicing generates two main forms, TFPI α and TFPI β (PMID: 26149025, PMID: 16246254). TFPI α is a soluble form of TFPI. It is a ~46 kDa glycoprotein and consists of an acidic N-terminal region followed by three tandem, Kunitz-type protease inhibitor domains and a basic C-terminal region. Based on protein mass, TFPI β (28 kDa) is considerably smaller than TFPI α (36 kDa), but both migrate with the same apparent molecular mass (46 kDa) on sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE), suggesting a difference in post-translational modifications (PMID: 16246254). TFPI associated with LDL is 34 kDa and lacks a substantial portion of the carboxy terminus including at least a portion of the third Kunitz domain. TFPI associated with HDL is 41 kDa and is in part a mixed disulphide complex between TFPI and apolipoprotein AII (PMID: 8979119). TFPI exerts its anticoagulant effects by inhibiting TF-FVIIa in a manner that is dependent on its inhibition of factor Xa (FXa) (PMID: 2927510).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31765-1-AP (TFPI antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.