For Research Use Only

E-Cadherin Polyclonal antibody

Catalog Number: 31863-1-AP



Purification Method:

WB 1:1000-1:4000 IF/ICC 1:200-1:800

Antigen affinity Purification

Recommended Dilutions:

Basic Information

Catalog Number: 31863-1-AP

Size:

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_009864.3

GeneID (NCBI):

150ul , Concentration: 550 ug/ml by

12550

Nanodrop: **UNIPROT ID:**

P09803 Full Name: cadherin 1

Calculated MW:

98kDa

Observed MW: 130 kDa

Applications

Tested Applications:

WB, IF/ICC, ELISA Species Specificity:

human, mouse

Positive Controls:

WB: MCF-7 cells, 4T1 cells

IF/ICC: MCF-7 cells,

Background Information

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. E-cadherin (epithelial cadherin), also known as CDH1 (cadherin 1) or CAM 120/80, is a classical member of the cadherin superfamily which also include N-, P-, R-, and B-cadherins. E-cadherin is expressed on the cell surface in most epithelial tissues. The extracellular region of E-cadherin establishes calcium-dependent homophilic trans binding, providing specific interaction with adjacent cells, while the cytoplasmic domain is connected to the actin cytoskeleton through the interaction with p120-, a-, β -, and γ -catenin (plakoglobin). E-cadherin is important in the maintenance of the epithelial integrity, and is involved in mechanisms regulating proliferation, differentiation, and survival of epithelial cell. E-cadherin may also play a role in tumorigenesis. It is considered to be an invasion suppressor protein and its loss is an indicator of high tumor aggressiveness. E-cadherin is sensitive to trypsin digestion in the absence of Ca2+.

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31863-1-AP (E-Cadherin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using E-Cadherin antibody (31863-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).