

For Research Use Only

SHB Polyclonal antibody

Catalog Number: 31967-1-AP



Basic Information

Catalog Number:

31967-1-AP

Size:

150ul , Concentration: 350 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG35936

GenBank Accession Number:

BC136581

GeneID (NCBI):

6461

UNIPROT ID:

Q15464

Full Name:

Src homology 2 domain containing adaptor protein B

Calculated MW:

55 kDa

Observed MW:

55 kDa

Purification Method:

Antigen affinity Purification

Recommended Dilutions:

WB 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : HUVEC cells, L02 cells, C2C12 cells, H9C2 cells

Background Information

SH2 domain-containing adapter protein B (SHB) is an adaptor protein that operates downstream of tyrosine kinases. It is crucial in various cellular responses, including cell survival, differentiation, and proliferation (PMID: 29792386). SHB is involved in the modulation of focal adhesion kinase (FAK) signaling, which is important for regulating cell cycle progression in hematopoietic stem cells (PMID: 23528453). The protein has also been implicated in the progression of myeloproliferative disorders by affecting the activity of FAK and other signaling intermediates (PMID: 24952416).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31967-1-AP (SHB antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.