

For Research Use Only

CD73 Polyclonal antibody

Catalog Number: 32131-1-AP



Basic Information

Catalog Number:

32131-1-AP

Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_011851.4

GeneID (NCBI):

23959

UNIPROT ID:

Q61503

Full Name:

5' nucleotidase, ecto

Calculated MW:

64kDa

Observed MW:

65-70 kDa

Purification Method:

Antigen affinity Purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

mouse

Positive Controls:

WB : mouse kidney tissue, mouse liver tissue, mouse lung tissue, mouse spleen tissue

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane-bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32131-1-AP (CD73 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.