For Research Use Only

CD147 Polyclonal antibody

Catalog Number: 32187-1-AP



Basic Information

Catalog Number: 32187-1-AP

Size:

GenBank Accession Number:

NM 009768.2

GeneID (NCBI):

150ul, Concentration: 500 ug/ml by 12215

Nanodrop: **UNIPROT ID:**

Source: P18572-1 Rabbit Full Name: Isotype: basigin IgG

Calculated MW: 42kDa

Observed MW: 35-55 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: RAW 264.7 cells, mouse small intestine tissue, mouse spleen tissue, mouse thymus tissue

Purification Method:

WB 1:5000-1:50000 IHC 1:500-1:2000

Antigen affinity Purification

Recommended Dilutions:

IHC: mouse small intestine tissue, mouse liver tissue

Background Information

CD147, also known as Basigin or extracellular matrix metalloproteinase inducer (EMMPRIN), is a transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 7812975). The molecule is composed of an intracellular portion, an extracellular portion, and a single transmembrane region. CD147 is expressed in various cell types (e.g., hematopoietic, epithelial, and endothelial cells) and at varying levels (PMID: 32968061). Increased expression of CD147 occurs in many tumors. CD147 is a pleiotropic molecule that plays an important role in fetal, neuronal, lymphocyte, and extracellular matrix development (PMID: 17945211). CD147 has been identified as a receptor essential for erythrocyte invasion by Plasmodium falciparum (PMID: 22080952). It has been reported thatthe spike protein of SARS-CoV-2 binds to CD147 on host cells, thereby mediating the viral invasion (Wang, Ke, et al, BioRxiv, 2020).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


*** 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32187-1-AP (CD147 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 32187-1-AP (CD147 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 32187-1-AP (CD147 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).