For Research Use Only

RBP4 Polyclonal antibody

Catalog Number: 32392-1-AP



Basic Information

Catalog Number:

32392-1-AP

Nanodrop:

Source:

Rabbit

Size:

GenBank Accession Number:

NM_011255.3

GeneID (NCBI):

150ul, Concentration: 450 ug/ml by 19662

UNIPROT ID: Q00724

Full Name: Isotype: retinol binding protein 4, plasma

IgG Calculated MW:

23kDa

Observed MW: 17-21 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

mouse, rat

Purification Method: Antigen affinity Purification Recommended Dilutions: WB 1:500-1:3000

Positive Controls:

WB: mouse heart tissue, mouse liver tissue, rat heart

tissue, rat liver tissue

Background Information

RBP4 (retinol-binding protein 4) is a carrier protein that transports vitamin A (retinol) from the liver to the peripheral tissues. Synthesized primarily by hepatocytes and adipocytes as a 21 kDa non-glycosylated protein, RBP4 is secreted into the circulation as a retinol-RBP4 complex. In plasma the RBP4-retinol complex is bound to transthyretin (TRR), which prevents prevent kidney filtration. Two truncated forms of RBP4, RBP4-L (truncated at Leu-183) and RBP4-LL (truncated at Leu-182 and Leu-183), exist by proteolytic process. RBP4-L and RBP4-LL, which do not bind TTR, are normally excreted into the urine but accumulate in the serum during renal failure. Urinary RBP4 has been reported as marker for glomerular disease. RBP4 also was identified as an adipokine that elevated in some INS-resistant states. Measurement of serum RBP4 could be used to assess the risk of INS resistance, type 2 diabetes, obesity, and cardiovascular disease. (18752671, 16034410)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32392-1-AP (RBP4 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.