

For Research Use Only

F8A2 Polyclonal antibody

Catalog Number:32696-1-AP



Basic Information

Catalog Number: 32696-1-AP	GenBank Accession Number: NM_012151.3	Purification Method: Antigen affinity Purification
Size: 150ul , Concentration: 300 ug/ml by Nanodrop;	GeneID (NCBI): 474383	Recommended Dilutions: WB: 1:500-1:2000 IF/ICC: 1:50-1:500
Source: Rabbit	UNIPROT ID: P23610	
Isotype: IgG	Full Name: coagulation factor VIII-associated (intronic transcript) 2	
Immunogen Catalog Number: AG38529	Calculated MW: 39kDa 371aa	
	Observed MW: 38 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: human, mouse, rat	WB : OVCAR-3 cells, SKOV-3 cells, mouse brain tissue, rat brain tissue
	IF/ICC : HeLa cells,

Background Information

F8A2 (Coagulation Factor VIII Associated 2) is a protein associated with coagulation factor VIII, which is widely expressed in a variety of cell types, localized mainly in the cytoplasm and nucleus, and also co-localized with Huntington's protein (HTT). Although its specific function is not fully defined, F8A2 may play an important role in intracellular and vesicular transport. In addition, F8A2 co-localizes with ubiquitin in neuronal cells and may be involved in the pathological process of neurodegenerative diseases.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32696-1-AP (F8A2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 32696-1-AP (F8A2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using F8A2 antibody (32696-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).