For Research Use Only

Carbonic Anhydrase 6/CA6 Polyclonal antibody

Catalog Number:32740-1-AP

Basic Information	Catalog Number: 32740-1-AP	GenBank Accession Number: NM_001215.3	Purification Method: Antigen affinity Purification
	Size: 150ul, Concentration: 450 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: EG2963	GeneID (NCBI): 765 UNIPROT ID: P23280-1 Full Name: carbonic anhydrase VI Calculated MW: 35 kDa Observed MW: 35-40 kDa	Recommended Dilutions: WB: 1:2000-1:10000
Applications	Tested Applications: WB, ELISA Species Specificity: human, pig	Positive Controls: WB : pig salivary gland tissue, human saliva	
Background Information	CA6 (Carbonic anhydrase VI) is a member of CA family proteins that play a central role in pH regulation and electrolyte balance. CA6 is also known as gustin, is a zinc-containing secreted protein which catalyzes the hydration of carbon hydroxide in saliva. CA6 is specifically expressed in the salivary gland of a number of mammalian species (PMID:28423504). The amino acid sequences are highly conserved across the species. And it was reported that decreasing of CA6 protein was associated with loss of taste and pathological morphology of taste buds (PMID: 9784398).		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	storage	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 32740-1-AP (Carbonic Anhydrase 6/CA6 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. Various lysates were subjected to SDS PAGE followed by western blot with 32740-1-AP (Carbonic Anhydrase 6/CA6 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.