

For Research Use Only

TAFA1/FAM19A1 Polyclonal antibody

Catalog Number:32791-1-AP



Basic Information

Catalog Number: 32791-1-AP	GenBank Accession Number: BC025746	Purification Method: Antigen affinity Purification
Size: 150ul , Concentration: 750 ug/ml by Nanodrop;	GeneID (NCBI): 407738	Recommended Dilutions: WB: 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q7Z5A9	
Isotype: IgG	Full Name: family with sequence similarity 19 (chemokine (C-C motif)-like), member A1	
Immunogen Catalog Number: AG37253	Observed MW: 15 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse brain tissue, rat brain tissue, rat cerebellum
Species Specificity: human, mouse, rat	

Background Information

TAFA1, also named as FAM19A1, is a secreted protein that belongs to a small family of cytokines characterized by a conserved FAFA (FAM19 and TAFA) structural domain. It is predominantly expressed in specific regions of the central nervous system, particularly the brain. Functionally, TAFA1 acts as a chemokine-like factor, playing a crucial role in regulating immune cell migration and activation within the brain, thereby contributing to neuroimmune communication. Furthermore, growing evidence highlights its significance in various pathological processes, including modulating neuroinflammation, influencing the progression of neurodegenerative diseases, and even exhibiting context-dependent roles in certain cancers. TAFA1/FAM19A1 can be detected as about 15 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32791-1-AP (TAF1/FAM19A1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.