

For Research Use Only

# CD158K/KIR3DL2 Polyclonal antibody, PBS Only

Catalog Number:33010-1-PBS



## Basic Information

<b>Catalog Number:</b> 33010-1-PBS	<b>GenBank Accession Number:</b> NM_006737.4	<b>Purification Method:</b> Antigen affinity Purification
<b>Size:</b> 100ug , Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 3812	
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P43630-1	
<b>Isotype:</b> IgG	<b>Full Name:</b> killer cell immunoglobulin-like receptor, three domains, long cytoplasmic tail, 2	
<b>Immunogen Catalog Number:</b> EG4355	<b>Calculated MW:</b> 50 kDa	
	<b>Observed MW:</b> 50-60 kda	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
human, mouse

## Background Information

KIR3DL2, also known as CD158k, is an inhibitory receptor expressed by natural killer (NK) cells and a subset of CD8+ T cells. It plays a role in immune regulation by binding to HLA class I molecules, specifically HLA-A3 and HLA-A11, in a peptide-dependent fashion. KIR3DL2 can also function as an innate immune receptor for delivering CpG DNA to TLR9 in NK cells, highlighting its role in both adaptive and innate immunity. This protein is primarily expressed on the cell surface, particularly in NK cells and certain T cells. It has been implicated in various diseases, particularly in the context of hematological malignancies and autoimmune diseases. Aberrant expression of KIR3DL2 has been observed in neoplastic cells in transformed mycosis fungoides and Sézary syndrome, suggesting its potential as a therapeutic target in primary cutaneous anaplastic large-cell lymphoma (pcALCL).


## Storage

**Storage:**  
Store at -80°C.  
**Storage Buffer:**  
PBS only, pH7.3

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 33010-1-AP (CD158K/KIR3DL2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 33010-1-PBS in a different storage buffer formulation.