

For Research Use Only

NKX2-4 Polyclonal antibody, PBS Only

Catalog Number: 33052-1-PBS



Basic Information

Catalog Number:

33052-1-PBS

Size:

100ug, Concentration: 1 mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG37291

GenBank Accession Number:

NM_033176

GeneID (NCBI):

644524

UNIPROT ID:

Q9H2Z4

Full Name:

NK2 homeobox 4

Calculated MW:

36kDa

Observed MW:

38-41 kDa

Purification Method:

Antigen affinity Purification

Applications

Tested Applications:

WB, IF/ICC, Indirect ELISA

Species Specificity:

human

Background Information

The NKX2-4 protein (full name: NK2 homeobox 4) belongs to the NK-2 type homeobox transcription factor family. Its main function is to act as a potential transcription factor, binding specifically to DNA sequences via its homeodomain and participating in the regulation of cell differentiation and the development of multicellular organisms. Studies have shown that this protein is active in the nucleus, and its aberrant expression may contribute to the development of acute myeloid leukemia by interfering with the normal differentiation of megakaryocytes and erythrocytes. Additionally, it can regulate the expression of genes such as TERT through allele-specific binding, thereby influencing tumor susceptibility in melanoma.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of NKX2-4 antibody (33052-1-AP; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-NKX2-4 transfected HEK-293 cells. This data was developed using the same antibody clone with 33052-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 33052-1-AP (NKX2-4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 33052-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using NKX2-4 antibody (33052-1-AP) at dilution of 1:200 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 33052-1-PBS in a different storage buffer formulation.