

For Research Use Only

C12orf43 Polyclonal antibody

Catalog Number: 33971-1-AP



Basic Information

Catalog Number: 33971-1-AP	GenBank Accession Number: BC014661	Purification Method: Antigen affinity Purification
Size: 150ul , Concentration: 530 ug/ml by Nanodrop;	GeneID (NCBI): 64897	Recommended Dilutions: WB: 1:500-1:2000
Source: Rabbit	UNIPROT ID: Q96C57	
Isotype: IgG	Full Name: chromosome 12 open reading frame 43	
Immunogen Catalog Number: AG40576	Observed MW: 37 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, LNCaP cells, ACHN cells, Calu-1 cells
Species Specificity: human	

Background Information

C12orf43 (also known as CUSTOS) is a nuclear-localized protein. It is predicted to be involved in the formation of the Spemann organizer, a key process in embryonic development, and acts as a negative regulator of the Wnt signaling pathway. Studies have shown that C12orf43 is differentially expressed in various tumors, and its high expression in liver cancer and lung cancer is significantly associated with poor patient prognosis, establishing it as an unfavorable prognostic marker. Additionally, C12orf43 has been found to be associated with neurodegenerative diseases, autoimmune disorders, and other conditions.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



WB result of C12orf43 antibody (33971-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-C12orf43 transfected LNCaP cells.



Various lysates were subjected to SDS PAGE followed by western blot with 33971-1-AP (C12orf43 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.